


Revision PPQu – Unit 3: Moles and Calculations

1. A student adds 2.12 g of sodium carbonate to an excess of dilute hydrochloric acid.

The chemical equation for the reaction is



Calculate the maximum mass, in g, of carbon dioxide formed in the reaction. (3)

mass = g

(Total marks for question = 3)

2. Compound Z contains 38.7% carbon, 9.7% hydrogen and 51.6% oxygen by mass.

(i) Show by calculation that the empirical formula of compound Z is CH_3O (2)

(ii) The relative formula mass (M_r) of compound Z is 62

Deduce the molecular formula of compound Z. (2)

molecular formula =

(Total for question = 4 marks)

3.

- (a) A student produces dry crystals of hydrated copper(II) sulfate from a saturated solution of copper(II) sulfate.
He calculates that 6.40 g of dry crystals should be formed.
The mass of dry crystals he actually obtains is 1.80 g less than he calculated.
Calculate the student's percentage yield.
Give your answer to one decimal place. (3)

percentage yield = %


- (b) Gypsum is hydrated calcium sulfate.
A sample of gypsum contains 79% of calcium sulfate by mass.
Calculate the value of x in $\text{CaSO}_4 \cdot x\text{H}_2\text{O}$
[M_r of CaSO_4 = 136 M_r of H_2O = 18] (3)

x =

(Total marks for question = 6)

4.

(a) The diagram shows the apparatus a teacher uses to determine the formula of an oxide of lead.



This is the teacher's method.

Step 1 find the mass of the reduction tube

Step 2 add some of the lead oxide to the reduction tube

Step 3 find the mass of the reduction tube and lead oxide

Step 4 pass hydrogen gas over the lead oxide and ignite the hydrogen at the hole

Step 5 heat the lead oxide strongly for 10 minutes

Step 6 keep passing hydrogen through the reduction tube until the tube and contents are cool

Step 7 find the new mass of the reduction tube and its contents

(i) Give a reason why hydrogen is passed through the reduction tube until the tube and contents are cool. (1)

.....
.....

(ii) Describe what the teacher should do next to make sure all the lead oxide has been reduced to lead. (2)

.....
.....
.....
.....

(b) The teacher completes the experiment and obtains these results.

mass of reduction tube = 23.50 g
mass of tube + lead oxide = 28.64 g
mass of tube + lead = 28.16 g

(i) Calculate the mass of lead formed. (1)

mass of lead = g

(ii) Calculate the mass of oxygen removed from the lead oxide. (1)


mass of oxygen = g

(iii) Determine the empirical formula of the lead oxide. (4)

empirical formula of the lead oxide

(c) The insoluble salt lead(II) chloride (PbCl_2) can be prepared by reacting a solution of lead(II) nitrate with dilute hydrochloric acid.

(i) Complete the equation for the reaction by adding the state symbols. (1)



(ii) Show that the maximum mass of lead(II) chloride that can be made from 0.0370 mol of hydrochloric acid is about 5 g.

$[M_r \text{ of } \text{PbCl}_2 = 278]$ (3)

maximum mass = g

(Total marks for question = 13)


TOTAL MARKS FOR QUESTIONS = 26

EXTENSIONS

1. A sample containing 0.02 moles of hydrated copper(II) sulfate is heated using a Bunsen burner.

The products of the reaction are anhydrous copper(II) sulfate and water.

This is the equation for the reaction:



Calculate the maximum number of water molecules in tube B after the sample of hydrated copper(II) sulfate has completely reacted.

One mole of any substance contains 6×10^{23} particles. (2)

maximum number of molecules =

2. The equation represents the complete combustion of an alkene.



Complete combustion of 0.0100 mol of the alkene produces 2.16 g of water.


Determine the molecular formula of this alkene.

[for H_2O , $M_r = 18$] (3)

molecular formula =

3. In experiment, a student adds 0.55 g of zinc to a solution containing 2.50×10^{-2} moles of hydrochloric acid.

Use the equation to show that hydrochloric acid is in excess.



[Ar of Zn = 65]

(2)

TOTAL MARKS FOR EXTENSION = 7