

Please check the examination details below before entering your candidate information

Candidate surname					Other names				
Centre Number					Candidate Number				

Pearson Edexcel International GCSE

Monday 18 November 2024

Morning (Time: 1 hour 15 minutes)

Paper reference **4CH1/2C**

Chemistry
UNIT: 4CH1
PAPER: 2C

You must have:
 Calculator

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 – *there may be more space than you need.*
- Show all the steps in any calculations and state the units.

Information

- The total mark for this paper is 70.
- The marks for **each** question are shown in brackets
 – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P75948A

©2024 Pearson Education Ltd.
 V:1/1/1/1/1/1/1/



2

A standard 1D barcode with vertical black bars of varying widths on a white background. Below the bars, the corresponding alphanumeric characters are printed: P 7 5 9 4 8 A 0 2 2 4.

Cu and Cl have not been rounded to the nearest whole number.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



Answer ALL questions.

Some questions must be answered with a cross in a box ☒. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☒.

- 1** The table shows the composition of five species that are either atoms or ions.

Species	Number of protons	Number of neutrons	Number of electrons
A	1	0	1
B	3	3	3
C	3	4	3
D	9	10	10
E	13	14	13

- (a) Use the table to answer these questions.

You may use each letter once, more than once, or not at all.

- (i) Give the letter of the species that is a negative ion.

(1)

- (ii) Give the letter of the species that has a full outer shell of electrons.

(1)

- (iii) Give the letter of the species that has an atomic number equal to its mass number.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (b) Explain why species B and C are atoms of the same element with identical chemical properties.

(2)

- (c) A proton has a mass of 1.6726×10^{-24} g.

A neutron has a mass of 1.6740×10^{-24} g.

Calculate the mass, in grams, of the nucleus of an atom of species C.

(2)

mass = g


(Total for Question 1 = 7 marks)



2 This question is about metals and metal compounds.

(a) All metals are malleable and good conductors of electricity.

The diagram shows the structure of a metal.




(i) Explain why metals are malleable.

(2)

(ii) Explain why metals are good conductors of electricity.

(2)

3 A teacher uses this apparatus to separate a mixture of ethanol and water.



(a) Give the name of this method of separation.

(1)

- ☐ **A** chromatography
- ☐ **B** crystallisation
- ☐ **C** filtration
- ☐ **D** fractional distillation

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(2)

.....

.....

(Total for Question 3 = 6 marks)



4 A teacher adds a small piece of lithium to water and collects the gas produced.

(a) (i) Give two observations when lithium is added to water.

(2)

1

.....

2

.....

(ii) The teacher adds a few drops of universal indicator to the solution at the end of the reaction.

Explain the final colour of the universal indicator.

(2)


.....

.....

.....

.....

(b) This is the equation for the reaction between lithium and water.



The teacher collects 550 cm^3 of hydrogen at rtp.

Calculate the mass, in grams, of lithium that the teacher added to the water.

[one mole of gas at rtp has a volume of $24\,000\text{ cm}^3$]

[for lithium, $A_r = 7$]

[for carbon $A_r = 12$ for hydrogen, $A_r = 1$ for oxygen $A_r = 16$]

(3)

mass = g



(c) The teacher then adds a small piece of potassium to water.

(i) Give one observation seen with potassium that is **not** seen with lithium.

(1)

(ii) Explain why potassium is more reactive than lithium.

Refer to atomic structure in your answer.

(3)

(Total for Question 4 = 11 marks)



- 5 A student does some titrations to find the volume of dilute nitric acid needed to exactly neutralise 25.0 cm^3 of sodium hydroxide solution.

This is the student's method.

Step 1 add 25.0 cm^3 of sodium hydroxide solution to a conical flask

Step 2 add three drops of methyl orange indicator

Step 3 fill a burette with the acid

Step 4 add acid from the burette until the indicator changes colour

Step 5 record the volume of acid added

- (a) (i) Give the name of the apparatus that the student should use to measure the volume of sodium hydroxide solution in step 1.

(1)

- (ii) Give the colour change seen in step 4.

(1)


- (iii) Give a reason why the student does not use universal indicator in this titration.

(1)


- (b) The student completes a rough titration and four accurate titrations.

The diagram shows the burette readings from the rough titration.

Start



End



The table shows the student's results.

	Rough titration	Titration 1	Titration 2	Titration 3	Titration 4
Burette reading at end in cm^3		21.80	22.85	21.75	24.10
Burette reading at start in cm^3		0.50	0.15	0.25	0.10
Volume added in cm^3		21.30	22.70	21.50	24.00

- (i) Complete the table by adding the results from the rough titration.

Record the volumes to the nearest 0.05 cm^3

(2)

- (ii) Concordant results are results within 0.20 cm^3 of each other.


Use the concordant results from the table to calculate the mean volume of acid added.

(3)

mean volume of acid = cm^3



- (c) This is the equation for the reaction between dilute nitric acid and sodium hydroxide solution.



After the titration, the student knows the volume of acid needed to neutralise 25.0 cm^3 of the sodium hydroxide solution.

Sodium nitrate decomposes at high temperatures.

Describe how the student could obtain pure, dry crystals of sodium nitrate from dilute nitric acid and sodium hydroxide solution.

(5)

(Total for Question 5 = 13 marks)



6 This question is about crude oil.

(a) Describe how crude oil is separated into fractions by fractional distillation.

(4)

(b) Some of the long-chain alkanes obtained from fractional distillation are cracked, producing shorter-chain alkanes and ethene.

(i) Give the conditions necessary for cracking to occur.

(2)


(ii) Explain why shorter-chain alkanes are more useful than longer-chain alkanes.

(2)



(iii) Ethene reacts with bromine.

The equation shows the displayed formulae of the reactants and product.



The table shows some bond energies.

Bond	Bond energy in kJ/mol
C=C	612
C—C	348
C—H	414
Br—Br	193
C—Br	276

Show that the molar enthalpy change, ΔH , for this reaction is about -100 kJ/mol.

(3)

$\Delta H =$ kJ/mol



(iv) Explain why the reaction between ethene and bromine is exothermic.

Refer to bond energies in your answer.

(2)

.....

.....

.....

.....

.....

.....

(Total for Question 6 = 13 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



7 This question is about alcohols and carboxylic acids.

(a) These are the structural formulae of alcohol A and carboxylic acid B.

Alcohol A $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$

Carboxylic acid B $\text{CH}_3(\text{CH}_2)_n\text{COOH}$

(i) Name alcohol A.

(1)

(ii) Draw the displayed formula of alcohol A.

(1)

(iii) Carboxylic acid B has a chain of carbon atoms with no branches.

The number of CH_2 units is represented by the letter n.

Calculate the value of n.

[for carboxylic acid B, $M_r = 242$]

(3)

n =

(iv) Alcohol A and carboxylic acid B react together to form an ester.

Give the other product of the reaction.

(1)




(b) These are the structural formulae of dicarboxylic acid C and diol D.

Dicarboxylic acid C $\text{HOOCCH}_2\text{CH}_2\text{COOH}$

Diol D $\text{HOCH}_2\text{CH}_2\text{OH}$

- (i) This is the equation for the reaction between dicarboxylic acid C and sodium hydroxide solution.



25.0 cm^3 of 0.150 mol/dm^3 sodium hydroxide solution is completely neutralised by 17.5 cm^3 of a solution of dicarboxylic acid C.

Calculate the concentration, in mol/dm^3 , of the solution of dicarboxylic acid C.

Give your answer to three significant figures.

(4)

concentration = mol/dm^3



(ii) Dicarboxylic acid C and diol D react to form a polyester.

Draw the displayed formula of the repeat unit of this polyester.

(2)

(Total for Question 7 = 12 marks)

TOTAL FOR PAPER = 70 MARKS



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE

